

Chemical Analysis of Irrigation Water and its Effects on Soil of Dera Ghazi Khan

Ayaz Ahmad^{1*}, Dr.Asma Saeed², Dr.Allah Bakhsh Gulshan³

1.*Assistant Professor of Biology Govt. Degree College Block.17 D.G.Khan,Punjab Pakistan

2. Associate Professor Department of Biological Sciences Gomal University D.I.K, KPK.
Pakistan.

3. Associate Professor of Botany,Registerar Ghazi University D.G.Khan,Punjab.Pakistan

Abstract

Irrigated water brings some dissolved salts from sources. The quality and quantity of these dissolved salts depend on the source. Usually most water dissolved substance include sodium (Na^{+1}), magnesium (Mg^{+2}), calcium (Ca^{+2}), chloride (Cl^{-1}), carbonate (CO_3^{-2}) and bicarbonates (HCO_3^{-1}). The amount and concentration of these dissolved ions determine the fitness of water for irrigation. The common quality parameters are Electrical Conductivity (EC), Sodium Adsorption Ratio (SAR) and Residual Sodium Carbonate (RSC). Unfortunately, canal water is inadequate to meet the requirement of crops under severe harvesting system in Punjab. In Dera Ghazi Khan there is many ways are used to irrigate the cultivated soils other than canal systems like tube wells. In which more than 70% water are of poor quality. The study area is either canal irrigated or irrigated through tube wells. This research was conducted in 25 km area from Qasba Samina Sadat to Taunsa canal west of the D.G.Khan city in 2018. To achieve the goal of this research, the study was revealed in the Soil and Water Testing Laboratory for Research Dera Ghazi Khan. The 40 water samples were collected from different sites which are irrigated through both canal and tube well system. The samples were collected in plastic bottles after the 30 minutes operation of tube wells. Similarly 22 soil samples were collected in plastic bags to test the effect of irrigated water on soil of the study area. Table.4 Shows that approximately 40% irrigated water is unfit and 20% is marginally fit for irrigation. This is indication of soil is towards salinity pattern. The water and soil samples were analyzed at Soil and Water Testing Laboratory for Research, Dera Ghazi Khan for electrical conductivity, cations (Ca^{+2} , Mg^{+2} , Na^{+1}) and anions (CO_3^{-2} , HCO_3^{-1} and Cl^{-1}) by the methods described by Page et al.,(1982)[10]. Residual sodium carbonates (RSC) and Sodium Adsorption Ratio (SAR) were calculated through international standard.

Key words. Irrigation water, EC, SAR, RSC, Salinity.

Introduction

Water is essential component of all forms of life and it is mainly obtained from two sources, i.e. surface water which includes rivers, canals, fresh water lakes, streams etc. and ground water like well water and borehole water [1]. Because of unique chemical properties of water due to its polarity and hydrogen bonding, it has ability to suspend, dissolve, absorb and adsorb many different compounds. Thus water is not pure in nature, as it acquires contaminants from its surrounding as well as other biological activities [2]. However Irrigated water brings some dissolved salts from sources [3]. The quality and quantity of these dissolved salts depend on the source. Usually most water dissolved substance include sodium (Na^{+1}), magnesium (Mg^{+2}), calcium (Ca^{+2}), chloride (Cl^{-1}), carbonate (CO_3^{-2}) and bicarbonates (HCO_3^{-1}). The amount and concentration of these dissolved ions determine the fitness of water for irrigation [4]. The common quality parameters are Electrical Conductivity (EC), Sodium Adsorption Ratio (SAR) and Residual Sodium Carbonate (RSC). Chemically irrigated water affect the soil by changing the nutrients composition in the soil [5]. Salts are originated from weathering of rocks and soil, including dissolution of lime, gypsum and other slowly dissolved soil minerals. These substances are transferred from water resource to wherever it is used [6]. Lower quality of irrigated water adversely affects the production rate of crops. But unfortunately quality of irrigated water should be deserted [7]. Unfortunately, canal water is inadequate to meet the requirement of crops under severe harvesting system in Punjab [8]. In Dera Ghazi Khan there is many ways are used to irrigate the cultivated soils other than canal systems like tube wells. In which more than 70% water are of poor quality [9]. Keeping in view the importance of irrigation water quality, the present study was contemplated with the following objectives

- i) To ensure the quality of tube wells water for irrigation and
- ii) Suggest the different option to protect the soil of district Dera Ghazi Khan from salination.

Materials and Methods

Description of study area

Dera Ghazi Khan District is situated between river Indus and Suleiman range and lies between 20.40 North and 70.75 East. The study area is either canal irrigated or irrigated through tube wells. This research was conducted in 25 km area from Qasba Samina Sadat to Taunsa canal west of the D.G.Khan city in 2018.

Water and soil sampling

To achieve the goal of this research, the study was revealed in the Soil and Water Testing Laboratory for Research Dera Ghazi Khan. The 40 water samples were collected from different sites which are irrigated through both canal and tube well system. The samples were collected in plastic bottles after the 30 minutes operation of tube wells. Similarly 22 soil samples were collected in plastic bags to test the effect of irrigated water on soil of the study area.

Sample analysis

The water and soil samples were analyzed at Soil and Water Testing Laboratory for Research, Dera Ghazi Khan for electrical conductivity, cations (Ca^{+2} , Mg^{+2} , Na^{+1}) and anions (CO_3^{-2} , HCO_3^{-1} and Cl^{-1}) by the methods described by Page et al.,(1982)[10]. Residual sodium carbonates (RSC) and Sodium Adsorption Ratio (SAR) were calculated through international standard. The collected water samples were analyzed on the basis of following criteria regarding their suitability for irrigation and effect of this water on salinization of soils irrigated by such water.

Results

Table.1 Criteria used for water analysis

| Parameters | Fit | Marginally Fit | Unfit |
|------------|--------|----------------|-------|
| EC us/cm | 0-100 | 100-1250 | >1250 |
| SAR | 0-6 | 6-10 | >10 |
| RSC meq/l | 0-1.25 | 1.25-2.50 | >2.50 |

Table.2 Chemical Analysis of Water samples

Total 40 irrigated water samples were analysed and their different parameters are as under

| Sample No. | EC us/cm | $\text{Ca}^{+}+\text{Mg}^{++}$ meq/l | Na^{+1} meq/l | CO_3^{-2} meq/l | HCO_3^{-1} meq/l | Cl^{-1} meq/l | SAR | RSC meq/l | Fit/unfit for irrigation |
|------------|----------|--------------------------------------|------------------------|--------------------------|---------------------------|------------------------|------|-----------|--------------------------|
| 1 | 616 | 5.78 | 0.38 | 3.21 | 3.97 | 1.5 | 0.22 | 1-23 | Fit |
| 2 | 2930 | 23.95 | 5.35 | 13.4 | 24.9 | 7 | 1.55 | 3.21 | Unfit |
| 3 | 1443 | 13.20 | 1.23 | 10.03 | 11.1 | 3 | 0.48 | 2.68 | Unfit |
| 4 | 948 | 7.37 | 2.11 | 6.43 | 7.3 | 2 | 1.1 | 1.21 | Fit |

| | | | | | | | | | |
|----|------|-------|-------|-------|------|-----|------|------|------------------|
| 5 | 418 | 3.84 | 0.34 | 3.1 | 2.9 | 1 | 0.25 | 1.04 | Fit |
| 6 | 1436 | 12.17 | 2.19 | 11.4 | 10.8 | 3 | 0.87 | 2.52 | Unfit |
| 7 | 1294 | 10.40 | 2.54 | 8.5 | 9.7 | 3 | 1.11 | 2.72 | Unfit |
| 8 | 647 | 6.05 | 0.42 | 4.21 | 5.03 | 1 | 0.24 | 1.16 | Fit |
| 9 | 1839 | 12.77 | 5.62 | 12.4 | 13.6 | 4 | 2.22 | 4.51 | Unfit |
| 10 | 272 | 2.65 | 0.07 | 1.2 | 1.5 | 1 | 0.06 | 0.28 | Fit |
| 11 | 276 | 2.37 | 0.39 | 1.4 | 1.5 | 1 | 0.36 | 0.25 | Fit |
| 12 | 2070 | 19.09 | 1.61 | 13.7 | 15.4 | 4.5 | 0.52 | 3.24 | Unfit |
| 13 | 588 | 4.81 | 1.07 | 3.5 | 4.6 | 1 | 0.69 | 1.20 | Fit |
| 14 | 1250 | 8.78 | 3.72 | 8.9 | 9.8 | 2.5 | 1.77 | 2.12 | M marginally Fit |
| 15 | 1145 | 6.87 | 4.58 | 7.6 | 8.7 | 2.5 | 2.48 | 2.04 | M marginally Fit |
| 16 | 4370 | 23.28 | 20.42 | 33.4 | 35.2 | 7 | 5.98 | 3.08 | Unfit |
| 17 | 504 | 3.89 | 1.15 | 2.5 | 3.3 | 1.5 | 0.83 | 0.37 | Fit |
| 18 | 1122 | 10.27 | 0.95 | 6.8 | 8.9 | 2 | 0.42 | 2.23 | M marginally Fit |
| 19 | 2480 | 23.91 | 0.89 | 12.86 | 18.7 | 5 | 0.26 | 2.85 | Unfit |
| 20 | 783 | 6.60 | 1.23 | 5.4 | 5.2 | 2 | 0.68 | 1.04 | Fit |
| 21 | 463 | 4.08 | 0.55 | 2.9 | 3.3 | 1 | 0.38 | 0.89 | Fit |
| 22 | 1754 | 11.57 | 5.97 | 11.9 | 13.5 | 3.5 | 2.48 | 2.25 | Fit |
| 23 | 2050 | 19.17 | 1.33 | 14.5 | 15.3 | 6 | 0.43 | 2.64 | Unfit |
| 24 | 960 | 8.51 | 1.09 | 4.5 | 6.7 | 3 | 0.53 | 0.85 | Fit |
| 25 | 2320 | 21.87 | 1.33 | 12.4 | 15.8 | 7 | 0.4 | 2.67 | Unfit |
| 26 | 1130 | 10.59 | 0.71 | 6.4 | 8.5 | 2.5 | 0.31 | 2.42 | M marginally Fit |
| 27 | 2080 | 19.49 | 1.31 | 14.5 | 16.1 | 4.5 | 0.42 | 2.82 | Unfit |
| 28 | 2090 | 19.41 | 1.49 | 14.2 | 15.4 | 5 | 0.48 | 2.92 | Unfit |
| 29 | 1016 | 9.29 | 0.87 | 5.4 | 7.2 | 2.5 | 0.40 | 1.48 | M marginally Fit |
| 30 | 990 | 9.12 | 0.78 | 6.2 | 7.1 | 2.5 | 0.36 | 0.89 | Fit |
| 31 | 2110 | 19.03 | 2.07 | 12.8 | 15.4 | 5.5 | 0.67 | 3.04 | Unfit |
| 32 | 1167 | 10.07 | 1.6 | 5.2 | 7.7 | 3.5 | 0.71 | 2.94 | M marginally |

| | | | | | | | | | Fit |
|----|------|-------|------|------|------|-----|------|------|------------------|
| 33 | 1146 | 10.02 | 1.44 | 6.8 | 7.18 | 3 | 0.64 | 2.23 | M marginally Fit |
| 34 | 998 | 9.03 | 0.95 | 6.03 | 7.05 | 2.5 | 0.45 | 1.04 | Fit |
| 35 | 1011 | 9.32 | 0.79 | 6.2 | 7.2 | 2.5 | 0.37 | 2.28 | M marginally Fit |
| 36 | 841 | 7.66 | 0.75 | 5.2 | 6.1 | 2 | 0.38 | 1.22 | Fit |
| 37 | 288 | 1.98 | 0.9 | 0.24 | 1.45 | 1 | 0.91 | 0.54 | Fit |
| 38 | 462 | 3.87 | 0.75 | 2.5 | 3.4 | 1 | 0.54 | 0.68 | Fit |
| 39 | 2050 | 19.01 | 1.49 | 12.5 | 14.8 | 5.5 | 0.48 | 2.21 | Unfit |
| 40 | 2060 | 20.01 | 1.79 | 14.5 | 15.6 | 5.7 | 0.60 | 2.68 | Unfit |

Table. 3 Ranges. Means and Standard Deviation of different Parameters

| | EC us/cm | Ca ⁺ +Mg ⁺⁺ meq/l | Na ⁺¹ meq/l | CO ₃ ⁻² meq/l | HCO ₃ ⁻¹ meq/l | Cl ⁻ meq/l | SAR | RSC meq/l |
|-----------------------|-------------|--|---------------------------|--|---|--------------------------|---------------|---------------|
| Range | 70.99 | 0.19-9.92 | 0.0- 9.90 | 0.02- 11.3 | 0.04- 9.90 | 0-9.69 | 0.02- 9.57 | 0.01- 10.3 |
| Mean | 1230.4 | 8.78 | 4.89 | 2.8 | 7.06 | 4.42 | 3.063 | 2.8 |
| Standard Deviation | 825.8 | 15.47 | 4.67 | 2.24 | 9.56 | 5.26 | 2.28 | 2.12 |

Table 4 Classification of water samples

| Total Samples | Fit for Irrigation | | Marginally fit for irrigation | | Unfit for irrigation | |
|------------------|--------------------------|------|--|------|----------------------|------|
| | Samples | %age | Samples | %age | Samples | %age |
| 40 | 16 | 40 | 08 | 20 | 16 | 40 |

Table 4. Shows that approximately 40% irrigated water is unfit and 20% is marginally fit for irrigation. This is indication of soil is towards salinity pattern.

Table 5 :Classification of water sample on the basis of EC(µS/m)

| Total Samples | Fit for Irrigation | | Marginally fit for irrigation | | Unfit for irrigation | |
|---------------|--------------------|------|-------------------------------|------|----------------------|------|
| | Samples | %age | Samples | %age | Samples | %age |
| 40 | 00 | 0.0 | 25 | 62.5 | 15 | 37.5 |

Table 6 :Classification of water sample on the basis of SAR

| Total Samples | Fit for Irrigation | | Marginally fit for irrigation | | Unfit for irrigation | |
|---------------|--------------------|------|-------------------------------|------|----------------------|------|
| | Samples | %age | Samples | %age | Samples | %age |
| 40 | 16 | 40 | 08 | 20 | 16 | 40 |

Table 7:Classification of water sample on the basis of RSC meq/l

| Total Samples | Fit for Irrigation | | Marginally fit for irrigation | | Unfit for irrigation | |
|---------------|--------------------|------|-------------------------------|------|----------------------|------|
| | Samples | %age | Samples | %age | Samples | %age |
| 40 | 20 | 50 | 07 | 17.5 | 13 | 32.5 |

Soil samples Analysis

During the current Study 22 soil samples were collected from the different sites which are irrigated with canal as well as tube well water. The analysis indicates that due to different salts in irrigation water all the selected soils were saline. The time will come very soon that these soils which are irrigated by canal as well as tube well water will be saline.

Table.1 Results of analysed soil samples

| Sample No. | EC ms/cm | Soil pH | Saturation % | Texture | Soil condition |
|------------|----------|---------|--------------|-----------|----------------|
| 1 | 4.22 | 8.13 | 50 | Clay loam | Saline |
| 2 | 11.70 | 7.48 | 52 | Clay loam | Saline |
| 3 | 11.18 | 7.57 | 18 | Sandy | Saline |
| 4 | 9.67 | 7.70 | 50 | Clay loam | Saline |

| | | | | | |
|----|-------|------|----|------------|--------|
| 5 | 28.80 | 7.50 | 36 | Loam | Saline |
| 6 | 5.04 | 8.07 | 52 | Clay loam | Saline |
| 7 | 14.50 | 7.46 | 56 | Clay loam | Saline |
| 8 | 7.93 | 7.98 | 54 | Clay loam | Saline |
| 9 | 15.92 | 7.11 | 50 | Clay loam | Saline |
| 10 | 14.79 | 7.54 | 48 | Clay loam | Saline |
| 11 | 25.90 | 7.96 | 56 | Clay loam | Saline |
| 12 | 15.98 | 7.98 | 28 | Sandy loam | Saline |
| 13 | 6.56 | 8.12 | 54 | Clay loam | Saline |
| 14 | 4.70 | 8.12 | 56 | Clay loam | Saline |
| 15 | 4.06 | 7.83 | 50 | Clay loam | Saline |
| 16 | 6.78 | 7.86 | 52 | Clay loam | Saline |
| 17 | 10.15 | 7.66 | 58 | Clay loam | Saline |
| 18 | 8.29 | 7.72 | 26 | Sandy loam | Saline |
| 19 | 9.76 | 7.62 | 48 | Clay loam | Saline |
| 20 | 13.75 | 7.42 | 52 | Clay loam | Saline |
| 21 | 8.41 | 7.83 | 54 | Clay loam | Saline |
| 22 | 7.42 | 7.4 | 26 | Sandy loam | Saline |

Table.1 of the soil samples indicates that due to high EC and pH values of the collected samples from the sites which are irrigated by canal water and tube wells are saline soils.

Discussion

In present study it is clearly indicated that in D.G.Khan the irrigation water is not fit for crop production in soils. The irrigated water from canals is no pure instead contains such substances which play main contribution in making the soil saline. Similarly tube well water is basically ground water, which is basically heavy water having many dissolved salts contribute in salinity.

Suggestions

Keeping in view the above results the farmers of this area are advised, before utilizing the water for irrigation, should be properly analysed from agriculture department. This is the only way to protect the soils from salinity.

References

- 1 .Afzal, S., I. Ahmad, M. Younas, M.D. Zahid, M.H.A. Khan, A. Ijaz and K. Ali. 2000. Study of water quality of Hudiara Drain, India-Pakistan. *Environ. Geol.* 26:87-96.
2. Afzal, S., M. Farooq, M.H.A. Khan, M.I. Khan and K. Ali. 1998. Nitrate, fecal coliform and pesticides in shallow groundwater of Raiwind, Lahore, Pakistan. *Proc. 67th Annual Session Symposium on Environment Protection and Resource Conservation by Pakistan Engineering Congress XXV*:14-22.
3. Ayers, R.S. and D.W. Westcot. 1985. Water quality for agriculture. *FAO Irri. Drain.* 29:1-109.
4. Barzegar, A.R., J.M. Oades, P. Rengasamy and L. Giles. 1994. Effect of sodicity and salinity on disaggregation and tensile strength of an alfisol under different cropping systems. *Soil Tillage Res.* 32:329-345.
5. Bhutta, M.N., M. Ramzan and C.A. Hafeez. 2002. Ground water quality and availability in Pakistan. *Proc. Seminar on "Strategies to Address the Present and Future Water Quality Issues"*. Pakistan Council of Research in Water Resources (PCRWR), Islamabad, Pakistan.
6. Chitmanat, C. and S. Traichaiyaporn. 2010. Spatial and temporal variations of physical-chemical water quality and some heavy metals in water, sediments and fishes of the Mae Kuang River, Northern Thailand. *Int. J. Agric. Biol.* 12:816-820.
7. Joshi, D.M., A. Kumar and N. Agrawal. 2009. Assessment of the irrigation water quality of River Ganga in Haridwar District India.
8. Malik, D.M., M.A. Khan and T.A. Chaudhry. 1984. Analysis method for soil, plant, and water. Soil Fertility Survey and Soil Testing Institute, Punjab, Lahore, Pakistan.
9. McMurry, J. and R.C. Fay. 2004. Hydrogen, Oxygen and Water, 4th Ed. In: K.P. Hamann (ed.). *McMurry Fay Chemistry*. Pearson Education, New Jersey. pp.575-599.
10. Mendie, U. 2005. The Nature of Water. In: *The Theory and Practice of Clean Water Production for Domestic and Industrial Use*. Lacto-Medals Publishers, Lagos. pp.1-21.