

PRILIMINARY PHYTOCHEMICAL SCREENING AND ANTIFUNGAL SUSCEPTIBILITY OF *SECURINEGA VIROSA* EXTRACT ON SOME UROPATHOGENS

Aernan Tracy Paulyn¹, Odo Joel Inya^{2*}, Zakari Ene Agnes¹

¹Departments of Microbiology, University of Agriculture, Makurdi, Nigeria

²Department of Fisheries and Aquaculture University of Agriculture, Makurdi, Nigeria

*Corresponding author:

odojoel@gmail.com

ABSTRACT

Securinega virosa is commonly called “Bush weed” and is one of the great African medicinal plants described as a true “cure all”. This study is aimed at evaluating the phytochemical screening and antifungal susceptibility of *Securinega virosa* extract on some uropathogen. Evidence of this study shows that *Securinega virosa* contains some phytochemicals such as alkaloids, saponin, glycosides, tannins, steroids, although, quinones and flavonoids are absent. Fungal isolates were obtained from the microbiology laboratory, Federal University of Agriculture, Makurdi. Susceptibility test was carried out using Agar well diffusion method. The MIC of the potent extract was determined according to the macro broth dilution technique. At the concentration of 200mg/ml, *Candida albicans* had the zone of inhibition of 9.00 ± 1.00 higher than the zone of inhibition of *Aspergillus niger*. The zone of inhibition of *Aspergillus niger* at 250mg/ml (12.92 ± 0.57) was higher than that of *Candida albicans* (12.33 ± 0.57). The potency of the aqueous leaf extract of *Securinega virosa* which reduced with decreased in concentration, 500mg/ml has the highest zone of inhibition across the isolates while 100mg/ml showed the lowest zone of inhibition. The study reveals that *S. virosa* show significant antifungal activities against *Candida* infection with a minimum inhibitory concentration of about 25%.

KEYWORDS: *Securinega virosa*, Susceptibility, *Candida albicans*, *Aspergillus niger*, Phytochemical, Uropathogen.

1. INTRODUCTION

Securinega virosa is one of the great African medicinal plants described as a true “cure all”, of which all parts are used as remedies, particularly the root (Neuwinger, 2016). It belongs to the family, Euphorbiaceae of the order, Geranaiales. *S. virosa* is a dense, low branching, with many branched shrub, sometimes a small spreading tree up to about 6 meters high, although, more commonly 2 to 3 meters, evergreen or deciduous. It is widely distributed throughout tropical Africa, also in India, Malaya, China and Australia (Dalziel, 2013).

In Nigeria, it is found in virtually all parts of the country. The common vernacular names of *S. virosa* include “*tsuwaawun karee, gussu, gwiiwar karee*” (Hausa), “*iranje*” (Yoruba), “*njisi nta*” (Ibo), “*shim shim*” (Kanuri), “*kartfi kartfi*” (Shuwa Arabs) and “*camal, cambe, came*” (Fulani). In many parts of Africa including the north Eastern Nigeria, the root and leafy twig decoctions are used for the treatment of epilepsy. The plant is said to have a hallucinogenic effect and the decoction of the root with other plant is used in Northern Nigeria for the treatment of mental illness (Neuwinger, 2016).

The plant is also known as common bush weed and rich in Antioxidants which protect the human body against free radicals that may cause pathological conditions. This plant (*Securinega virosa*) produces definite phytochemicals with physiological actions in the human body. The medicinal values of the plants lie in the component of their phytochemicals. Phytochemicals are secondary metabolites produced by plants (Hill, 2010). Research interest has been directed to the use of natural phytochemical for the formulation of antioxidant-based drugs (Mondon *et al.*, 2012). The majority of the active antioxidant compounds are flavonoids, flavones, isoflavones, anthocyanins, cumarins, lignans, catachins, phenols and isocatachins. Flavonoids and flavones which are some of the constituent of the plant are widely distributed secondary metabolites with antioxidant and antiradical properties (Augustin *et al.*, 2015). The leaves are used in the treatment of fever, body pain, stomachache, rheumatism, diarrhea, pneumonia, diabetes and epilepsy.

Among the most common infectious diseases, urinary tract infections (UTIs) are a commonly encountered diseases by clinicians in developing countries with an estimated annual global incidence of at least 250 million (Ronald *et al.*, 2001). UTIs refer to the presence of microbial pathogens within the urinary tract and it is usually classified by the infection site:-bladder (cystitis), kidney (pyelonephritis), or urine (bacteriuria) and also can be asymptomatic or symptomatic, UTIs that occur in a normal genitourinary tract with no prior instrumentation are considered as “uncomplicated,” whereas “complicated” infections are diagnosed in genitourinary tracts that have structural or functional abnormalities, including instrumentation such as indwelling urethral catheters, and are frequently asymptomatic (Gonzalez and Schaeffer, 1999). It has been estimated that globally symptomatic UTIs result in as many as 7 million visits to outpatient clinics, 1 million visits to emergency departments, and 100,000 hospitalizations annually (Wilson and Gaido, 2004).

The aim of this study is to evaluate the preliminary phytochemical screening and antifungal susceptibility of *Securinega virosa* extracts on some uropathogens.

3.0 MATERIALS AND METHODS

3.1 Study Area

The plant sample (*Securinega virosa*) was collected from Federal University of Agriculture, Makurdi, Benue state. The plant sample was then taken to microbiology research laboratory and identified by a botanist before analysis.

3.2 Equipment/ Reagents

The equipment and reagents are as follows, Text tube rack, Conical Flask, digital weighing balance, Pipette, mortar, Heating mantle, concentrated sulphuric acid, glacial acetic acid, dilute hydrochloric acid, distilled water, Acetic anhydride, Ferric Chloride, distilled water and Methanol.

3.3 Preparation of plant sample

The samples (*Securinega virosa*) were washed under running water, dried for 3 days under room temperature and grinded into powdery form using mortar and pestle. The grinded sample was sieved and stored in an airtight bottle.

3.3.1 Extraction

180g of the plant sample was weighted and soaked in 500ml of methanol and water respectively. The soaked sample was allowed to stand for 2 days (48hr), it was filtered and concentrated in a water bath at 49.7°C

3.3.2 Procedure for Phytochemical Screening

Major metabolites classes such as alkaloids, glycosides, flavonoids, Tannins saponins, steroids, Quinones, protein, phenol and terpenoids were screened according to the methods described by a guide to modern techniques of plant analysis, Medicinal Plants and Traditional Medicine in Africa and Pharmacognosy

3.3.3 Test for Phenol

2ml of the extract was pipette into a test tube, few drops of dilute ferric chloride solution was added. The formation of a red, blue, green, or purple coloration indicates the presence of phenols.

3.3.4 Test for Alkaloids

2ml of the extract was pipette into a test tube the filtrate was carefully tested with Mayer's reagent (potassium mercuric iodide). yellow coloured precipitate indicates the presence of alkaloids.

3.3.5 Test for Flavonoids

To 2ml of extract, few drops ammonia solution was added. Appearance of yellow or orange colour indicates the presence of flavonoids.

3.3.6 Test for Tannins

To 2 ml water extract of all plant parts, 2 ml of 10% ferric chloride solution was added in a test tube. Blue-black precipitate indicates the presence of tannins.

3.3.7 Test for Saponin

To 2ml methanolic extract of the plant, 2 ml distilled water was added in a test tube and vigorously shaken. Persistent froth volume produced, checked each 10 minutes for 30 minutes indicates the presence of saponin.

3.3.8 Test for Steroids

1g of the plant extract was dissolved in a few drops of acetic acid. It was gently warmed and cooled under the tap water and a drop of concentrated H_2SO_4 acid was added along the side of the test tube. The appearance of green colour indicates the presence of steroid.

3.3.9 Test of Terpenoids

Few drops of chloroform were reacted with 2ml of the latterly and few drops of concentrated H_2SO_4 was added. A reddish-brown precipitate produced immediately indicate the presence of terpenoids

3.4 ANTI-MICROBIAL ACTIVITY OF THE PALNT EXTRACT

3.4.1 Preparation of test organisms

Cold stored agar slant cultures of Gram positive and Gram-negative organism and fungi were used in this study. The fungal isolates were obtained from the microbiology laboratory, University of Agriculture Makurdi.

Viability test of each organism was carried out by resuscitating the organisms in buffered peptone both and there after sub-cultured into nutrient agar medium and incubated at 37°C for 24hrs. The probable identity of the clinically sourced isolates was further confirmed by exposing the cultures to an array of biochemical tests which includes coagulase, indole methylised, citrate, action and oxidase tests as described by Collins *et al* 2004 and Sharma (2009). The result of the biochemical reaction elicited by the test isolate were compared with standard identification keys as described by Collins *et al.*, 2004 on Sabouraud dextrose agar to check the purity.

3.4.2 Preparation of Concentration of the Plant Extracts.

One gram (1g) each of the aqueous and methanolic extracts was added to 2ml of distilled water and methanolic respectively to give a concentration of 250mg/ml, 100mg/ml and 58mg/ml were prepared by double broth dilution method as described by Udochukwu *et al.* (2015).

3.4.3 Anti-Microbial Susceptibility testing of the extract with the test organisms.

ANTI FUNGAL ASSAY

Susceptibility testing was carried out using agar well diffusion method. In this method, the inocula were prepared by inoculating the test organisms in dextrose broth and allowed to stand for 72hours. The cultures were diluted to 0.5 Mcfarland turbidity standards. About 0.5ml each of the cultured fungi was pipette into Petri dish after which prepared Sabouraud dextrose agar was pour plated and allowed to solidify.

After the cultured plates have gelled, wells were bored on the surface of the agar plates using 4mm core borer. About 0.2ml of the different concentration of each extract were transferred into the well using pasture pipette. The wells were sufficiently spaced to prevent the resulting zone of inhibition from overlapping. Fluconazole (25mg) was used as positive control. The plates were allowed to stand for 24 hours. The experiment was performed in triplicates and resulting zone of inhibition measured at the diameter of the well using a ruler.

Minimum Inhibitory Concentration (MIC)

The MIC of the potent extracts was determined according to the macro broth dilution technique described by Baron and Finegold (1990). Standardized suspension of the test organism was inoculated into a series of five sterile test tubes of nutrient broth containing two-fold dilution of the extracts and incubated at 37% for 24hours. The MIC was recorded as the least concentration that inhibited the growth of test organism.

3.5 Statistical Analysis

Data were analyzed for mean and standard deviation. Difference in parameter was tested for statistical difference at $p<0.05$ using T-test. All the analysis was done using statistical package service solution (SPSS) version 21.

RESULTS

4.1 Phytochemical Screening

Table 1 indicates the phytochemical constituents of each extract of *Securinega virosa* for the presence of secondary metabolites. Quinones and Flavonoid are absent in both methanolic and aqueous leaf extract of *Securinega virosa*.

Zone of inhibition of methanol leaf extract of *Securinega virosa* is presented on Table 2. *Candida albicans* have the highest zone of inhibition at the concentration of 500mg/ml which is significantly higher than the control. At the concentration of 200mg/ml, *Candida albicans* has the zone of inhibition of 9.00 ± 1.00 higher than the zone of inhibition of *Aspergillus niger*. The zone of inhibition of *Aspergillus niger* at 250mg/ml (12.92 ± 0.57) is higher than that of *Candida albicans* (12.33 ± 0.57). There was however no significant difference in the mean zone of inhibition at different concentrations of methanolic leaf extracts of *Securinega virosa* at $P > 0.05$.

Table 3 shows the potency of the aqueous leaf extract of *Securinega virosa* which reduced with decrease in concentration. 500mg/ml has the highest zone of inhibition across the isolates while 100mg/ml showed the lowest zone of inhibition. *Aspergillus niger* has the highest zone of inhibition across all concentrations except at 100mg/ml (9.33 ± 1.55 mm). There was significant different in the mean of inhibition at different concentrations of aqueous leaf extract of *Securinega virosa* at $P \leq 0.05$

Table 1: phytochemical properties of *Securinega virosa* leaf

| Phytochemical | Methanol | Aqueous |
|---------------|----------|---------|
| Glycosides | + | + |
| Steroids | + | + |
| Saponins | + | + |
| Quinones | - | - |
| Phenol | + | + |
| Terpenoids | + | + |
| Protein | + | + |
| Alkaloids | + | + |
| Flavonoid | - | - |
| Tannin | + | + |

Key:

(+) = present

(-) = Absent

Table 2: Zone of inhibition of methanol leaf extract of *Securinega virosa* on selected organisms

| Test organism | zone of inhibition (mm) | | | | | |
|--------------------|--------------------------|-------------|-------------|-------------|-------------|------------|
| | Control (Fluconazole) | 500mg/ml | 250mg/ml | 200mg/ml | 125mg/ml | 100mg/ml |
| <i>C. albicans</i> | 1.20±12.16 | 17.33±2.082 | 12.33±0.577 | 9.00±1.00 | 1.33±0.577 | 11.00±0.00 |
| <i>A. niger</i> | 1.15±11.03 | 16.00±2.000 | 12.92±1.564 | 12.33±1.528 | 10.33±0.577 | 9.00±1000 |
| (p> 0.05) Df=5 | | | | | | p=0.182 |

Table 3: Zone of inhibition of aqueous leaf extract of *Securinega virosa* on selected organisms

| Test organism | zone of inhibition (mm) | | | | | |
|-------------------|--------------------------|-------------|-------------|-------------|-------------|------------|
| | Control (Fluconazole) | 500mg/ml | 250mg/ml | 200mg/ml | 125mg/ml | 100mg/ml |
| <i>C. alblica</i> | 1.33±7.60 | 13.33±2.309 | 11.33±0.577 | 10.33±0.577 | 10.33±0.577 | 11.00±0.00 |
| <i>A. niger</i> | .88±11.87 | 15.67±1.528 | 14.33±1.528 | 12.33±1.528 | 10.33±1.528 | 9.33±1.55 |
| (p≤ 0.05) Df=5 | | | | | | p=0.026 |

5.1 DISCUSSION

Evidence of the study indicated that *Securinega virosa* plant contains phytochemicals such as alkaloid, saponin, tannin, glycoside although, quinones and flavanoids are absent. These phytochemicals are found in the leaf of the plant in various concentrations. This phytochemical gives the plant the therapeutic capacity which makes them a good medicinal source. The finding of the study also corroborates with the finding of Kiran Kumar *et al.* (2013) who also reported the presence of alkaloids, flavanoids, phenols, glycosides, tannins, saponins and lignins in the leaf extract of *Mirabilis jalapa*.

The relative susceptibility of *Candida albicans* to antibiotics is dependents on the age of culture because the culture environment is rapidly changing and the cell population become more physiologically heterogenous and hence, more resistant with age. So, it is beneficial to test the effect of the plant extract on *Candida albicans* culture in its stationary phase of growth.

The study of biological and chemical properties showed that the Nigerian flora has a real therapeutic and nutritional potential and can be used to treat or prevent many diseases. These plants have a real interest in human health for their antibacterial, Antifungal, antiplasmoidal and antioxidant activities.

The antifungal activity of the extracts of leaves of *S. virosa* was evaluated on two species of fungi which are: *C. albicans*, *Aspergillus niger*. These saprophytes and opportunistic yeast are present in the mucosa where they take advantage of an imbalance of existing flora or immune deficiency to multiply and cause infections. Resistance to azoles, including fluconazole, may explain clinical failures (Vanessa, 2001). Determining the diameter of the inhibition zones and determination of antifungal parameters (MIC) were used to assess their sensitivity. In view of the results, we note that the best extraction yields are obtained with the methanol extracts, as well as the best antifungal activity is obtained with ethanol extracts.

The methanol extracts were the most active; indicating that they concentrate most of the compounds. Methanol would be the best extraction solvent of active substances from this plant. It may also be explained that the activity of antibiotics in plant extracts against fungi or growth may be due to their mechanism of action, chemical structure or spectrum of activity.

As *S. virosa* induces diameters of inhibition zone greater than 10 mm and the MIC range from 100mg/ml. These results suggest that the plant has an antifungal activity. This corroborates with the results of Agassounon *et al.* (2001) which have shown that this plant has fungicidal properties.

The diameters of inhibition zones are 17 mm for methanolic extract the most active extract and 13 mm for aqueous extract at a concentration of 500 mg/ml. Kubmarawa *et al.* (2007) showed that the methanol extract of the leaves of *S. virosa* has significant inhibitory activity against *C. albicans* and *Aspergillus niger*. These results are supported by the work of Agassounon *et al.* (2001) which showed that the methanolic extract concentrate is better than aqueous extract of the leaf of plant. It should be noted that the antimicrobial activity of medicinal plants varies from one geographic region to another.

5.2 CONCLUSION

Study of *Securinega virosa* showed significant antifungi activities against multidrug resistant fungi that complicate *Candida* infection with an MIC of about 25%

Evidence from this study clearly shows that *Securinega virosa* contains phytochemicals that are essential for both the plant and makes the plant a good source of medicine for man. At 500mg/ml *Securinega virosa* extract has a minimum fungicidal concentration of 50%

5.3 RECOMMENDATION

From the result of this study, it is recommended that:

- i. Further studies should be carried out on the quantitative and molecular analysis of organisms associated with urinary tract infection.
- ii. In-vivo study of the plant extract should be carried out to determine its toxicity on human organ such as the bladder.
- iii. Research should be carried out on the antiviral effect of *Securinega virosa* extract on some selected organism particularly Hepatitis and rubella diseases.

REFERENCE

- [1] Agassounon DTM, De Souza C, Anani KT, Koumaglo K, Toukourou F, Gbeassor M, (2001);
a. Evaluation of the cytotoxic, antiviral, antibacterial and antifungal activities of six medicinal plants. *Pharm. Méd. Trad. AF JOOL*. 11:93-105.
- [2] Augustin F., Valerie P, Jacques F, and Virginie RV, (2015), Synthesis and biological evaluation of new securinine analogues as potential anticancer agents. *European Journal of Medicinal Chemistry*, 109: 287-293.
- [3] Baron, J.E. and Feingold, S.M. (1990). Methods of testing antimicrobial effectiveness. In Bailey Scotts diagnostic microbiology. Mosby C.V (Ed) Missouri, 171-194.
- [4] Collins, C.H., Lyne, P.M., Grange, J.M. and Falkinham, J.O. (2004). Collins and Lyne's Microbiological Methods, 8th Ed. Arnold Publication London, 465.
- [5] Dalziel A., (2013), Chemical composition of the fruit of tetraptera and the physicochemical properties of its oil. *Global J. Pure and Applied Science*. 3:61-67.
- [6] Gonzalez CM, Schaeffer AJ. (1999): Treatment of urinary tract infection: what's old, what's new, and what works. *World Journal Urol*. 17(6):372-82. doi: 10.1007/s003450050163. PMID:10654368
- [7] Hill V., (2010), Antibacterial activity of plant extracts and phytochemicals on antibiotic resistant Bacteria. *Brazilian journal of Microbiology*.31:247-256.
- [8] Kiran Kumar Angadi, Ammani Kandru and Abdul Rahaman (2013); Antihyperglycaemic, Antihyperlipidaemic and Antioxidant Assays (In Vivo) Of *Nymphaea Pubescens* Leaf Extract.
- [9] International Journal of Pharma and Bio Sciences; 4(2), 624 – 630, ISSN NO: 0975-6299
- [10] Kubmarawa D., G. A. Ajoku, N. M. Enwerem and D. A. Okorie (2007) Preliminary phytochemical and antimicrobial screening of 50 medicinal plants from Nigeria African Journal of Biotechnology Vol. 6 (14), pp. 1690-1696, <http://www.academicjournals.org/AJB>
- [11] Michael Wilson, Loretta Gaido (2004) Laboratory Diagnosis of Urinary Tract Infections in Adult a. Patients *Clinical Infectious Diseases*, Volume 38, Issue 8, 15 April 2004, Pages 1150–1158, <https://doi.org/10.1086/383029>
- [12] Mondon J., Todokoro, T. and Maekawa, A., (2012), Nutritional evaluation of chemical component of leaves and stems of sweet potatoes (ipomoea batatas L.). *Food Chem*. 68:359-367.
- [13] Neuwinger O., (2016), Proximate and mineral composition of the leaves of hairy indigo (*Indigofera astragalina*). *Pakistan Journal of Nutrition*.10(2):168-175.
- [14] Ronald AR, Nicolle LE, Stamm E, Krieger J, Warren J, Schaeffer A, Naber KG, Hooton TM, Johnson J, and Chambers S, Andriole V. (2001): Urinary tract infection in adults: research priorities and strategies. *International Journal of Antimicrobial Agents*. 17(4):343-8. doi: 10.1016/s0924-8579(01)00303-x. PMID: 11295419.
- [15] Sharma, K. (2009). *Manual of Microbiology* Ane Books. Pot. Limited. New Delhi, 405.

- [25] Udochukwu, U., Omoje, F.I., Uloma, I.S. and Oseiwe, I.D. (2015). Phytochemical analysis of *Vernonia amygdalina* and *Occimum gratissimum* extracts and their antimicrobial activities on some drug resistance bacteria. *American Journal of Research Communication*, 3(5) 225-235.
- [26] Vanessa C. (2001) Vaginal candidiasis recurrent to *Candida albicans*: Diagnosis, pathophysiology, treatment. University Henri Poincaré -Nancy 1, France. Thesis Pharmacy.; 131
- [27] pathophysiology, treatment. University Henri Poincaré -Nancy 1, France. Thesis Pharmacy.; 131